

# FT-IR and FT Raman: Experimental and Theoretical Studies and HOMO-LUMO Analysis of 1,2,4,5-Tetracyanobenzene

Ali Sanda Bawa<sup>1,\*</sup>, Abdoul-Rachid Chaibou Yacouba<sup>1</sup>, Seydou Ouedraogo<sup>2</sup>,  
Mabinty-Bayo Bangoura<sup>2</sup>, Karifa Bayo<sup>2</sup>, Marcel Bouvet<sup>3</sup>

<sup>1</sup>Department of Chemistry, University of Agadez, Agadez, Niger

<sup>2</sup>Department of Chemistry, University Ouaga I Pr. Joseph KI-ZERBO, Ouagadougou, Burkina Faso

<sup>3</sup>Institut de Chimie Moléculaire, University of Burgundy, Dijon, France

## Email address:

sandabawaali@yahoo.fr (Ali Sanda Bawa)

\*Corresponding author

## To cite this article:

Ali Sanda Bawa, Abdoul-Rachid Chaibou Yacouba, Seydou Ouedraogo, Mabinty-Bayo Bangoura, Karifa Bayo, Marcel Bouvet. FT-IR and FT Raman: Experimental and Theoretical Studies and HOMO-LUMO Analysis of 1,2,4,5-Tetracyanobenzene. *Science Journal of Chemistry*. Vol. 11, No. 4, 2023, pp. 161-167. doi: 10.11648/j.sjc.20231104.14

Received: July 11, 2023; Accepted: August 14, 2023; Published: August 31, 2023

**Abstract:** Experimental and theoretical studies on molecular and vibrational structure of 1,2,4,5-Tetracyanobenzene (TCNB) have been undertaken in the present work. The FT-IR spectrum and FT Raman spectrum of TCNB were recorded respectively in the region 4000–400  $\text{cm}^{-1}$  and 2500–400  $\text{cm}^{-1}$ . The optimized geometries in the ground state were calculated by DFT (B3LYP) methods with 6-311G basis set. The optimized structural shown the difference in the bond lengths and some angles are very little affected with a small reduction. The computational results indicate that the molecule adopts a plane geometry and belongs to the  $D_{2h}$  point group with the aromatic ring lying in the plane of symmetry. The molecular electrostatic potential shown that the two protons of TCNB as well as the benzene ring are in electron deficient region whereas the CN group are in electron rich region. The carbon atom of CN group and the carbon on the benzene ring are in antibonding interactions in HOMO and in bonding interactions in the LUMO. The harmonic vibrational frequencies, infrared intensities and Raman activities of the TCNB were evaluated. After scaling, the computational wavenumbers are in agreement with the experimental values. A detailed interpretation of the infrared and Raman spectra of TCNB is presented.

**Keywords:** Tetra-Cyanobenzene, DFT Calculations, Symmetry, IR and Raman

## 1. Introduction

When van der Waals interactions rule the building of organic molecular materials, their optical and electronic properties are logically related to the energy of the frontier molecular orbital, the highest occupied molecular orbital, HOMO, and the lowest unoccupied molecular orbital, LUMO, of the molecular units [1]. It is result an occupied binding molecular through the hybridization of the HOMO and LUMO orbital which causes a filled valence band in the solid state and makes the material semiconducting or insulating [2]. When strong intermolecular interactions, like donor–acceptor tandems or hydrogen-bonding, takes place in the solid state, material display generally unique crystalline structure and optoelectronic properties. The interaction between an electron

donating precursors with an electron accepting can produce a new molecular material called Charge Transfer Complexes (CTCs). Various molecules organics like 1,2,4,5-tetracyanobenzene (TCNB) can behave as electron acceptor and form molecular complexes with electron-donor molecules. In the molecule of 1,2,4,5-tetracyanobenzene, the two hydrogen atom as well the benzene ring are highly electron deficient due to the strong electron with-drawing effect of the four CN group. TCNB has been widely used to form CTCs in the solid state with different electron donors because of his important role as an electron acceptor [3-7]. Different vibrational studies have been reported on the TCNB molecule. For example, the calculations of the atomic movements of the fundamental vibrations of the TCNB were carried out in terms of the Cartesian displacement coordinates

and the precise vibrational modes were schematically drawn in diagrams [8]. The polarized IR and far-IR spectra of single crystals of TCNB were recorded and the normal coordinate calculated with the modified Urey-Bradley force field and the valence force field for the in-plane and out-of-plane vibrations, respectively [9]. Using electron diffraction and by ab initio calculations at different levels of theory, the molecular structure of TCNB in gas-phase has been determined [10]. Chakraborty and al. have characterized the vibrational levels of TCNB in the electronic ground and first excited states by dispersed fluorescence and laser induced fluorescence spectroscopy [11]. In this present work, the number of vibrational mode of TCNB in solid state was determined according to group theory and symmetry and the complete vibrational analysis was realized via the combination of the experimental and theoretical results obtained by density functional theory (DFT) based on scaled quantum chemical. Density functional theory calculations are reported to provide excellent vibrational frequencies of organic compounds, if the calculated frequencies are scaled to compensate for the approximate treatment of electron correlation, for basis set deficiencies and for the anharmonicity effects [12]. The vibrational wavenumbers, complete geometrical parameters, modes of vibrations and other parameters of TCNB molecule were investigated using B3LYP calculations with 6-311G basis sets. Specific scale factors were employed in the predicted frequencies for the accuracy.

## 2. Material and Methods

### 2.1. Experimental Details

The compound which is of spectroscopic grade was purchased from commercial sources and hence used directly for recording the spectra without further purification.

FT-IR spectrum were recorded between 4000 and 400 $\text{cm}^{-1}$  using a PerkinElmer Spectrum 100 FT-IR spectrometer. Raman spectroscopy was conducted employing a Renishaw in Via Raman microscope with a 633 nm laser between 400 and 2500  $\text{cm}^{-1}$ . The frequencies of all sharp bands are accurate to  $\pm 2 \text{ cm}^{-1}$ .

### 2.2. Theoretical Details

The molecular structure optimization of the TCNB molecule and corresponding vibrational harmonic frequencies were calculated using the DFT at Beckee-3-Lee-Yag-Parr (B3LYP) with 6-311G basis set using Gaussian 09W program package without any constraint on the geometry of the molecule [13]. The computed vibrational frequencies have been scaled in order to allow direct comparison with the experimental spectra. It is possible to obtain a better agreement between the computed and experimental frequencies by using appropriate scale factors for the different regions of vibrations. Vibrational frequency assignments were made with accurately by combining the results of the GAUSSVIEW 3 program with symmetry considerations [14-15].

## 3. Results and Discussion

### 3.1. Molecular Geometry and Parameters

The optimized molecular structures with the numeration of atoms and the choice of axes obtained from Gaussian 09 and GAUSSVIEW programs is as shown in Figure 1. The global minimum energy obtained by DFT structure optimization using 6-311G basis set for the title molecule as -601.157386332 u. a. The optimized structural parameters, such as bond lengths, bond angles, and dihedral angles determined at the same basis set are listed in Table 1.

The carbon-carbon bonds in TCNB are not in the same order of length as in the case of benzene. It appears a little distorted with shorter C<sub>1</sub>-C<sub>2</sub>, C<sub>2</sub>-C<sub>3</sub>, C<sub>4</sub>-C<sub>5</sub> and C<sub>5</sub>-C<sub>6</sub> and larger C<sub>1</sub>-C<sub>6</sub>, C<sub>3</sub>-C<sub>4</sub>. The larger bonds are about 1.4173 Å (141.73 pm) in length, whereas the shorter other are about 1.4 Å (140 pm) long. The difference in the bond lengths between the larger and shorter bonds is about 0.0173 Å. The difference of 0.02 Å value has been obtained at MP2 level using 6-31++G\*\* basis set [11].

In the C-C ring of the TCNB molecule, all carbons are hybridized sp<sup>2</sup>. This implies that the bond angles around the carbon will be exactly 120°. The computed results shown that the bonds angles slightly deviate from the ideal sp<sup>2</sup> hybridized values. Some angles are very little affected with a small reduction but this change give rises to a slight opening of the others angles. However, the computational results indicate that the molecule adopts a plane geometry and belongs to the D<sub>2h</sub> point group with the aromatic ring lying in the plane of symmetry.





Figure 1. Molecular structure of TCNB with numbering of atoms.

### 3.2. Frontier Molecular Orbital Analysis

The frontier molecular orbital (FMOs) play an important role in the optical and electric properties, as well as in quantum chemistry and UV-visible spectra. The HOMO-LUMO energy gap is an important measure of the stability of the compound. The HOMO has the ability to give an electron, thereby the energy of the molecule is corresponding to the ionization potential whereas the LUMO obtains an electron as an electron acceptor and it corresponds

to the electron affinity. Molecule with large values of frontier orbital gap, are less polarizable, with small chemical reactivity and high kinetic stability and is called a hard molecule and vice versa [16]. In our calculations the energy band gap of TCNB is calculated as 4.7737 eV. The Figure 2 shows the electron density of the HOMO and LUMO using density functional theory (DFT) at B3LYP/6-311G level for the optimized geometry in the ground state.

As seen from the orbital plots in Figure 2, there are antibonding interactions between the carbon atom of CN group and its direct neighbor on the benzene ring in HOMO while they are in bonding interactions in the LUMO.




**Figure 2.** Orbital plots showing the HOMO–LUMO electronic transitions in the TCNB.

### 3.3. Molecular Electrostatic Potential

The molecular electrostatic potential (MEP) has been particularly useful in depicting the molecular size and shape as

an indicator of the charge distributions of molecules. The MEP of the TCNB molecule showed in Figure 3 was calculated at the DFT/B3LYP/6–311G level of theory. The different color in the plot indicates the electrostatic potential regions in which the yellow-coloured region is nucleophile and electron rich, the blue colour indicates the electron-poor electrophile region and the white region indicates neutral atoms. As seen from the MEP, the two protons attached to carbons atom number 2 and 5 as well as the benzene ring are in electron deficient region whereas the CN groups are in electron rich region.



**Figure 3.** The molecular electrostatic potential surface of the TCNB calculated at the DFT/B3LYP/6-311G.

**Table 1.** Geometrical parameters optimized in 1,2,4,5-tetracyanobenzene, bond length (Å), bond angle (°) and dihedral angle (°).

| ParametersB3LYP/6-311G |          |              |          |              |          |
|------------------------|----------|--------------|----------|--------------|----------|
| Bond length (Å)        |          |              |          |              |          |
| C1-C2                  | 1.4      | C3-C4        | 1.4173   | C6-C13       | 1.4294   |
| C1-C6                  | 1.4173   | C3-C9        | 1.4294   | C9-N10       | 1.1663   |
| C1-C15                 | 1.4294   | C4-C5        | 1.4      | C11-N12      | 1.1663   |
| C2-C3                  | 1.4      | C4-C11       | 1.4294   | C13-N14      | 1.1663   |
| C2-H7                  | 1.0791   | C5-C6        | 1.4      |              |          |
| C15-N16                | 1.1663   | C5-H8        | 1.0791   |              |          |
| Bond angle (°)         |          |              |          |              |          |
| C2-C1-C6               | 119.5182 | C2-C3-C4     | 119.5182 | C4-C5-C6     | 120.9635 |
| C2-C1-C15              | 119.4869 | C2-C3-C9     | 119.4869 | C4-C5-H8     | 119.5182 |
| C6-C1-C15              | 120.9949 | C4-C3-C9     | 120.9949 | C6-C5-H8     | 119.5182 |
| C1-C2-C3               | 120.9635 | C3-C4-C5     | 119.5182 | C1-C6-C5     | 119.5182 |
| C1-C2-H7               | 119.5182 | C3-C4-C11    | 120.9949 | C1-C6-C13    | 120.9949 |
| C3-C2-H7               | 119.5182 | C5-C4-C11    | 119.4869 | C5-C6-C13    | 119.4869 |
| Dihedral angle (°)     |          |              |          |              |          |
| C6-C1-C2-C3            | 0.0      | C1-C2-C3-C4  | 0.0      | C3-C4-C5-C6  | 0.0      |
| C6-C1-C2-H7            | 180.0    | C1-C2-C3-C9  | 180.0    | C3-C4-C5-H8  | 180.0    |
| C15-C1-C2-C3           | 180.0    | H7-C2-C3-C4  | 180.0    | C11-C4-C5-C6 | 180.0    |
| C15-C1-C2-H7           | 0.0      | H7-C2-H3-C9  | 0.0      | C11-C4-C5-H8 | 0.0      |
| C2-C1-C6-C5            | 0.0      | C2-C3-C4-C5  | 0.0      | C4-C5-C6-C1  | 0.0      |
| C2-C1-C6-C13           | 180.0    | C2-C3-C4-C11 | 180.0    | C4-C5-C6-C13 | 180.0    |
| C15-C1-C6-C5           | 180.0    | C9-C3-C4-C5  | 180.0    | H8-C5-C6-C1  | 180.0    |
| C15-C1-C6-C13          | 0.0      | C9-C3-C4-C11 | 0.0      | H8-C5-C6-C13 | 0.0      |

### 3.4. Vibrational Modes

The maximum number of potentially active observable fundamentals modes of vibration from an on-linear molecule

which contains N atoms is equal to (3N-6), apart from three translational and three rotational degrees of freedom. The molecule of TCNB has 16 atoms and belongs to point group  $D_{2h}$ . According to the group theory, the three Cartesian

displacements of the 16 atoms provide 48 (3N) internal modes.

The species of fundamentals modes of vibrations are determined by considering the character of a reducible representation of dimension 3N by calculating the character of

the matrix representing the effect of each operation of symmetry on the base. Table 2 shows the character tables of point group  $D_{2h}$  and table 3 list the character of reducible representation  $\Gamma_{3N}$  for the same point group.

**Table 2.** Character table for the  $D_{2h}$  point group.

| $D_{2h}$ | E | $C_2(z)$ | $C_2(y)$ | $C_2(x)$ | i  | $\sigma_{(xy)}$ | $\sigma_{(xz)}$ | $\sigma_{(yz)}$ |                 |    |
|----------|---|----------|----------|----------|----|-----------------|-----------------|-----------------|-----------------|----|
| $A_g$    | 1 | 1        | 1        | 1        | 1  | 1               | 1               | 1               | $x^2, y^2, z^2$ |    |
| $B_{1g}$ | 1 | 1        | -1       | -1       | 1  | 1               | -1              | -1              | Rz              | xy |
| $B_{2g}$ | 1 | -1       | 1        | -1       | 1  | -1              | 1               | -1              | Ry              | xz |
| $B_{3g}$ | 1 | -1       | -1       | 1        | 1  | -1              | -1              | 1               | Rx              | yz |
| $A_u$    | 1 | 1        | 1        | 1        | -1 | -1              | -1              | -1              |                 |    |
| $B_{1u}$ | 1 | 1        | -1       | -1       | -1 | -1              | 1               | 1               | z               |    |
| $B_{2u}$ | 1 | -1       | 1        | -1       | -1 | 1               | -1              | 1               | y               |    |
| $B_{3u}$ | 1 | -1       | -1       | 1        | -1 | 1               | 1               | -1              | x               |    |

**Table 3.** Character of reducible representation of point group  $D_{2h}$ .

| $D_{2h}$                             | E  | $C_2(z)$ | $C_2(y)$ | $C_2(x)$ | i  | $\sigma_{(xy)}$ | $\sigma_{(xz)}$ | $\sigma_{(yz)}$ |
|--------------------------------------|----|----------|----------|----------|----|-----------------|-----------------|-----------------|
| Nj                                   | 16 | 4        | 0        | 0        | 0  | 0               | 4               | 16              |
| $2\cos\phi \pm 1$                    | 3  | -1       | -1       | -1       | -3 | 1               | 1               | 1               |
| $\Gamma_{3N} = Nj (2\cos\phi \pm 1)$ | 48 | -4       | 0        | 0        | 0  | 0               | 4               | 16              |

The reducible representation of dimension 3N was reduced using the formula of reduction:

$$a_i = \frac{1}{h} \sum_k \chi_i(R_k) * \chi_i(R_k) * n(R_k)$$

$h$ : is total number of operations in the point group;

$\chi_i(R_k)$ : character of the reducible representation of the operations for the equivalence class;

$\chi_i(R_k)$ : character of the irreducible representation of the operations for the equivalence class;

$n(R_k)$ : number of operations in the  $k^{\text{th}}$  column of the character table.

For the TCNB molecule,  $\Gamma_{3N} = 8A_g + 3B_{1g} + 5B_{2g} + 8B_{3g} + 3A_u + 8B_{1u} + 8B_{2u} + 5B_{3u}$

From the character table  $\Gamma_{trans} = B_{1u} + B_{2u} + B_{3u}$  and  $\Gamma_{rot} = B_{1g} + B_{2g} + B_{3g}$

Consequently by subtraction of the translation and rotation species, the complete set of vibrational modes is given by:

$$\Gamma_{vib} = 8A_g + 2B_{1g} + 4B_{2g} + 7B_{3g} + 3A_u + 7B_{1u} + 7B_{2u} + 4B_{3u}$$

The TCNB has forty two non-degenerate vibrations. For an N-atomic molecule, 2N-3 of all vibration is in-plane and N-3 is out-of-plane [17]. Thus twenty-nine of all the forty two vibrations are in plane and thirteen are out-of-plane. Since the molecule belongs to  $D_{2h}$  group, all vibrations being

symmetric through mirror plane of molecule  $\sigma_{(yz)} \equiv \sigma_h$  will belong to  $A_g$ ,  $B_{3g}$ ,  $B_{1u}$  and  $B_{2u}$  species and the others being asymmetric through  $\sigma_{(yz)} \equiv \sigma_h$  will belong to  $B_{1g}$ ,  $B_{2g}$ ,  $A_u$  and  $B_{3u}$  species.

The character table shows that  $B_{1u}$ ,  $B_{2u}$ ,  $B_{3u}$  are infrared active while  $A_g$ ,  $B_{1g}$ ,  $B_{2g}$  and  $B_{3g}$  are Raman active. Due to the center of inversion, the modes that are active in the IR are not active in the Raman, and vice versa. This is known as the mutual exclusion rule. Thus, IR and Raman spectra should show respectively 18 bands and 21 bands.

The detailed analysis of fundamental modes of vibration with FT-IR and FT Raman experimental frequencies scaled vibrational frequencies and IR intensity of TCNB using B3LYP method with 6-311G basis set is reported in Table 4. Sometimes, it is necessary to scale down the calculated harmonic frequencies in order to improve the calculated values in agreement with the experimental values. The scaled calculated frequencies minimize the root-mean square difference between calculated and experimental frequencies for bands with definite identifications. Hence, the harmonic calculated frequencies, infrared, and Raman intensities are scaled by 0.965 for the low frequency region and 0.986 for the high frequency region.

The observed experimental spectra; FT-IR recorded between 4000 and 400  $\text{cm}^{-1}$  and FT Raman recorded between 2500 and 400  $\text{cm}^{-1}$  compare to and theoretical spectra are showed respectively in Figures 4 and 5.

**Table 4.** Comparison of the observed and calculated vibrational frequencies 1,2,4,5-tetracyanobenzene.

| Mode | Species  | Experimental |       | TheoreticalB3LYP6-311G |              |        | Assignments   |
|------|----------|--------------|-------|------------------------|--------------|--------|---|
|      |          | FT-IR        | Raman | Frequency unscaled     | IR Intensity | Scaled |   |
| 1    | $A_u$    |              |       | 60.70                  | 0.0000       | 58.58  | $\gamma\text{C}-\text{CN}$                            |
| 2    | $B_{3u}$ |              |       | 96.48                  | 17.7608      | 93.11  | $\gamma\text{C}-\text{CN}$                            |
| 3    | $A_g$    |              |       | 114.91                 | 0.0000       | 110.89 | $\beta\text{C}-\text{CN}+\beta\text{C}\equiv\text{N}$ |
| 4    | $B_{2u}$ |              |       | 123.15                 | 2.0166       | 118.84 | $\beta\text{C}\equiv\text{N}+\beta\text{C}-\text{CN}$ |
| 5    | $B_{1u}$ |              |       | 130.06                 | 13.0501      | 125.51 | $\beta\text{C}-\text{CN}+\beta\text{C}\equiv\text{N}$ |

| Mode | Species         | Experimental |        | TheoreticalB3LYP6-311G |              |         | Assignments   |
|------|-----------------|--------------|--------|------------------------|--------------|---------|---|
|      |                 | FT-IR        | Raman  | Frequency unscaled     | IR Intensity | Scaled  |   |
| 6    | B <sub>2g</sub> |              |        | 158.51                 | 0.0000       | 152.96  | $\gamma\text{C}\equiv\text{N}$ , sym  |
| 7    | B <sub>1g</sub> |              |        | 219.70                 | 0.0000       | 212.01  | $\gamma\text{C}-\text{CN}$  |
| 8    | B <sub>3g</sub> |              |        | 233.95                 | 0.0000       | 225.76  | $\beta\text{C}\equiv\text{N}$   |
| 9    | A <sub>u</sub>  |              |        | 356.95                 | 0.0000       | 344.46  | $\gamma\text{C}-\text{CN}$  |
| 10   | B <sub>3g</sub> |              |        | 366.58                 | 0.0000       | 353.75  | $\nu\text{C}-\text{CN}$   |
| 11   | B <sub>3u</sub> |              |        | 387.49                 | 0.2418       | 373.92  | $\gamma\text{C}-\text{H}$   |
| 12   | A <sub>g</sub>  |              | 405w   | 416.92                 | 0.0000       | 402.33  | $\beta\text{C}-\text{CH}-\text{C}$ (ring) $\nu\text{C}-\text{CN}$             |
| 13   | B <sub>2g</sub> |              | 460vw  | 481.84                 | 0.0000       | 464.97  | $\gamma\text{C}-\text{C}\equiv\text{N}$                                       |
| 14   | B <sub>1u</sub> | 465w         |        | 494.11                 | 0.6543       | 476.82  | $\beta\text{C}-\text{C}\equiv\text{N}$  |
| 15   | B <sub>2u</sub> |              |        | 511.63                 | 1.1966       | 493.72  | $\beta\text{C}-\text{CN}$   |
| 16   | A <sub>g</sub>  |              | 506vw  | 528.90                 | 0.0000       | 510.39  | $\beta\text{C}\equiv\text{N}$ , $\beta\text{C}-\text{CN}$                     |
| 17   | B <sub>3u</sub> | 507vs        |        | 553.37                 | 9.9941       | 534.00  | $\gamma\text{CN}-\text{C}-\text{CH}-\text{C}-\text{CN}$                       |
| 18   | B <sub>1g</sub> |              | 550m   | 598.88                 | 0.0000       | 577.91  | $\gamma\text{C}-\text{CN}$  |
| 19   | B <sub>1u</sub> | 628vw        |        | 642.63                 | 3.4624       | 620.14  | Ring deformation  |
| 20   | B <sub>3g</sub> |              | 686vw  | 725.38                 | 0.0000       | 699.99  | $\beta\text{C}-\text{CN}$   |
| 21   | A <sub>g</sub>  |              | 723m   | 739.31                 | 0.0000       | 713.43  | $\beta\text{C}-\text{CH}-\text{C}$ ring<br>$\nu\text{CCN}-\text{CCN}$ ring    |
| 22   | A <sub>u</sub>  |              |        | 762.42                 | 0.0000       | 735.74  | $\gamma\text{C}-\text{CN}$ ring<br>$\gamma\text{C}-\text{C}$ ring deformation |
| 23   | B <sub>2u</sub> | 759w         |        | 782.06                 | 6.1802       | 754.69  | $\nu\text{C}-\text{CN}$ stretch   |
| 24   | B <sub>2g</sub> |              |        | 804.45                 | 0.0000       | 776.30  | $\gamma\text{C}-\text{H}$ sym, $\gamma\text{C}-\text{C}$ ring                 |
| 25   | B <sub>3u</sub> | 917vs        |        | 966.93                 | 41.6692      | 933.09  | $\gamma\text{C}-\text{H}$ asym  |
| 26   | B <sub>2g</sub> |              |        | 968.26                 | 0.0000       | 934.37  | $\gamma\text{C}-\text{H}$ sym   |
| 27   | B <sub>3g</sub> |              | 1034vw | 1063.38                | 0.0000       | 1026.16 | $\nu\text{C}-\text{CN}$ stretch<br>$\beta\text{C}-\text{C}$ ring              |
| 28   | B <sub>1u</sub> |              |        | 1208.89                | 3.7500       | 1166.58 | $\beta\text{C}-\text{CH}-\text{C}$<br>$\beta\text{C}-\text{C}-\text{CH}$      |
| 29   | B <sub>2u</sub> | 1182vw       |        | 1221.42                | 0.0366       | 1178.67 | $\beta\text{C}-\text{H}$  |
| 30   | A <sub>g</sub>  |              |        | 1276.91                | 0.0000       | 1232.22 | CCN-CH breathing  |
| 31   | B <sub>2u</sub> | 1277s        | 1261s  | 1308.26                | 12.0440      | 1262.47 | $\nu\text{CCN}-\text{CCN}$ (ring)<br>$\nu\text{CCN}-\text{CH}$ (ring)         |
| 32   | B <sub>3g</sub> |              |        | 1321.37                | 0.0000       | 1275.13 | $\beta\text{C}-\text{H}$  |
| 33   | B <sub>1u</sub> | 1387w        |        | 1401.72                | 3.1162       | 1352.66 | $\nu\text{CCN}-\text{CH}$ (ring)  |
| 34   | B <sub>2u</sub> | 1485s        |        | 1532.34                | 36.8901      | 1478.71 | $\beta\text{C}-\text{H}$  |
| 35   | A <sub>g</sub>  |              | 1540m  | 1550.99                | 0.0000       | 1496.71 | $\nu\text{CCN}-\text{CCN}$ (ring)   |
| 36   | B <sub>3g</sub> |              | 1598w  | 1636.94                | 0.0000       | 1579.64 | $\nu\text{CCN}-\text{CH}$ (ring)  |
| 37   | A <sub>g</sub>  |              | 2239vs | 2278.26                | 0.0000       | 2246.36 | $\nu\text{C}\equiv\text{N}$ stretch   |
| 38   | B <sub>3g</sub> |              | 2249sh | 2279.35                | 0.0000       | 2247.44 | $\nu\text{C}\equiv\text{N}$ stretch   |
| 39   | B <sub>2u</sub> | 2245s        |        | 2280.30                | 2.0959       | 2248.37 | $\nu\text{C}\equiv\text{N}$ stretch   |
| 40   | B <sub>1u</sub> |              |        | 2283.98                | 1.3649       | 2252.01 | $\nu\text{C}\equiv\text{N}$ stretch   |
| 41   | B <sub>1u</sub> | 3048s        |        | 3219.60                | 0.5509       | 3174.53 | $\nu\text{C}-\text{H}$ asym   |
| 42   | A <sub>g</sub>  | 3113s        |        | 3221.07                | 0.0000       | 3175.97 | $\nu\text{C}-\text{H}$ sym  |

w-weak, vw-veryweak, s-strong, vs-very strong, m-medium, sh-shoulder, v-stretching,  $\nu_{\text{asy}}$ -asymmetric stretching,  $\nu_{\text{sym}}$ -symmetric stretching,  $\beta$ -in plane bending,  $\gamma$ -out of plane bending,  $\omega$ -wagging, t-twisting,  $\delta$ -scissoring,  $\tau$ -torsion, r-rocking.




Figure 4. Comparison of experimental (a) and theoretical FT-IR spectra of TCNB.






Figure 5. Comparison of experimental and theoretical FT Raman spectra of TCNB.

#### C-H vibrations

The carbon-hydrogen vibrations in aromatic compounds can be found in the range  $3080\text{--}3030\text{ cm}^{-1}$  (C-H stretching) [18]. The corresponding C-H in plane and out of plane bending vibrations are usually observed between  $1000$  and  $1300\text{ cm}^{-1}$  and between  $750$  and  $1000\text{ cm}^{-1}$  respectively [19-20]. TCNB molecule has two C-H bonds which undergo two stretching vibrations associated with the ring. In the experimental IR spectrum of our title molecule, the asymmetric and symmetric C-H stretching are respectively observed as a fine and intense band at  $3048$  and  $3113\text{ cm}^{-1}$ . The assigned values are in good coherence with the results in the literature [9] and the B3LYP/6-311G basis set. In the experimental IR spectrum of benzene an intense band attributed to C-H stretching is observed at  $3064\text{ cm}^{-1}$  and at  $3036\text{ cm}^{-1}$  respectively in gas phase [21] and in a condensed phase [22].

In the calculated IR spectrum of our title molecule, the out of plane bending vibration  $\gamma$  (CH) are predicted at  $374\text{ cm}^{-1}$ ,  $776\text{ cm}^{-1}$ ,  $933\text{ cm}^{-1}$  and  $934\text{ cm}^{-1}$ . In the experimental IR spectrum; the very strong band  $917\text{ cm}^{-1}$  is attributed to the bending vibration out of plane of the C-H bond. The in plane bending vibrations,  $\beta$  (CH) are predicted in the calculated IR spectrum at  $1179\text{ cm}^{-1}$ ,  $1275\text{ cm}^{-1}$ ,  $1479\text{ cm}^{-1}$ . In the experimental IR spectrum, the corresponding bands are observed at  $1182\text{ cm}^{-1}$ ,  $1277\text{ cm}^{-1}$  and  $1485\text{ cm}^{-1}$  demonstrating a perfect agreement with the theory.

#### C-C vibrations

The ring carbon-carbon stretching vibrations in benzene derivatives usually occur in the region  $1400\text{--}1625\text{ cm}^{-1}$  [23]. In general, the bands are of variable intensity are observed at  $1256$ ,  $1605$ ,  $1388$ ,  $1278\text{ cm}^{-1}$  from the frequencies given in literature [24]. In this present work, the frequency observed at FT-IR and FT Raman spectra at  $1598$ ,  $1540$ ,  $1387$  are assigned to C=C stretching vibrations and the frequencies assigned at  $1261\text{ cm}^{-1}$  for C-C stretching vibrations. The theoretically computed frequencies values at  $1579\text{ cm}^{-1}$ ,  $1496\text{ cm}^{-1}$ ,  $1352\text{ cm}^{-1}$  and  $1262\text{ cm}^{-1}$  for C-C stretching vibrations by B3LYP 6-311G method are in excellent agreement with the experimental data.

#### Cyano group (C≡N) vibrations

For cyano compounds the  $\text{--C}\equiv\text{N}$  stretching absorption is observed in the narrow region between  $2260$  and  $2200\text{ cm}^{-1}$  [25]. It is frequent to observe the reduction of frequency to  $2240\text{--}2220\text{ cm}^{-1}$  when the cyano group is engaged in conjugation with aromatic ring. The  $\text{--C}\equiv\text{N}$  stretching band may be of variable intensity in infrared, and in Raman the band is medium to very strong intensity [26]. The very strong band at  $2239\text{ cm}^{-1}$  in Raman and a strong band at  $2245\text{ cm}^{-1}$  in IR is assigned to the  $\text{--C}\equiv\text{N}$  stretching of TCNB. The theoretically computed and assigned by Gaussview program at  $2246/2248\text{ cm}^{-1}$  is assigned to  $\text{--C}\equiv\text{N}$  stretching vibrations.

The in-plane bending of  $\text{--C}\equiv\text{N}$  bond is obtained in the low frequency region at  $507\text{ cm}^{-1}$  in infrared as strong and at  $506\text{ cm}^{-1}$  in the Raman as very weak. In the calculated spectrum; this band appears at  $510\text{ cm}^{-1}$ . All these assignments coincide well with the literature data and calculated scaled values [25].

## 4. Conclusion

We have carried out DFT calculations on the structure and vibrational spectra of TCNB. The vibrational frequencies analysis by B3LYP/6-311G method agree overall with experimental results. On the basis of agreement between the calculated and experimental results, assignments of all the fundamental vibrational modes of TCNB were examined and proposed. From the geometrical discussion, it was concluded that the substitutions of H atom by the  $\text{--CN}$  groups modify the  $\text{C}_1\text{--C}_6$ ,  $\text{C}_3\text{--C}_4$  bonds lengths thus the hexagonal nature of the benzene ring. The frequencies of vibration calculated by B3LYP 6-311G method are in excellent agreement with the experimental data.

## Acknowledgements

All authors thank sincerely, Prof Marcel BOUVET of University of Burgundy (ICMUB) for his availability.

## References

- [1] Ali Sanda Bawa, Rita Meunier-Prest, Yoann Rousselin, Jean-Pierre Couvercelle, Christine Stern, Bernard Malézieux and Marcel Bouvet; Series of charge transfer complexes obtained as crystals in a confined environment; *Cryst Eng Comm*, 2021, 23, 6418–6426.: 10.1039/D1CE00929J.
- [2] Micheal Huth, Organic Charge Transfert Systems: The next step in Molecular Electronics, Beilstein Bozen on Molecular Engineering and Control, Prien (Chiemsee), Germany 2012. <http://www.beilstein-institut.de/Bozen2012/Proceedings/Huth/Huth.pdf>
- [3] J. Fünfschilling, M. Samoc, D. F. Williams, Quenching of the delayed fluorescence by charge carriers in crystals of anthracene-TCNB CT complex; *Chem. Phys. Lett.* 96 (1983) 157-160. [https://doi.org/10.1016/0009-2614\(83\)80483-7](https://doi.org/10.1016/0009-2614(83)80483-7).
- [4] T. Kobayashi, S. Nagakura, Two-photon excitation inorganic crystals by dye laser; *Chem. Phys. Lett.* 13 (1972) 217. [https://doi.org/10.1016/0009-2614\(72\)85045-0](https://doi.org/10.1016/0009-2614(72)85045-0)
- [5] G. Agostini, C. Coravaja, G. Giacometti, L. Pasimeni, D. A. Clemente, Photoexcited triplets and structural properties of single crystals of the trans-stilbene-1,2,4,5-tetracyanobenzene (1: 2) complex, *J. Phys. Chem.* 92 (1988) 997. DOI: 10.1021/j100315a028.
- [6] H. Jiang, P. Hu, J. Ye, K. K. Zhang, Y. Long, W. Hu and C. kloc, *J. Mater. Chem. C*, 2018, 1884-1902.
- [7] X. Ye, Y. Liu, Q. Guo, Q. Han, C Ge; S Cui, L Zhang and X. Tao, *Nat. Commun.*, 2019, 10, 1-9.
- [8] J. Umemura and T. Takenaka, Cartesian Displacements of Normal Vibrations of 1,2,4,5-Tetracyanobenzene and 1,2,4,5-Tetracyanobenzene-d<sub>2</sub> Molecules, *Bull. Inst. Chem. Res., Kyoto Univ.*, 51, 206 (1973). [https://repository.kulib.kyoto-u.ac.jp/dspace/bitstream/2433/76783/1/chd056\\_4\\_176.pdf](https://repository.kulib.kyoto-u.ac.jp/dspace/bitstream/2433/76783/1/chd056_4_176.pdf)
- [9] Tohru TAKENAKA, Junzo UMEMURA, Shinichi TADOKORO, Shinzaburo OKA and Takashi KOBAYASH; Vibrational Spectra of 1,2,4,5-Tetracyanobenzene and 1,2,4,5-Tetracyanobenzene-d<sub>2</sub> Crystals; *Bull. Inst. Chem. Res., Kyoto Univ.*, Vol. 56, No. 4, 1978.
- [10] Schultz, G., Szabados, Á., Tarczay, G. *et al.* Molecular Structure of 1,2,4,5-Tetracyanobenzene from Gas-Phase Electron Diffraction and Theoretical Calculations. *Structural Chemistry* 10, 149–155 (1999). <https://doi.org/10.1023/A:1022089415252>
- [11] S. Chakraborty, S. Wategaonkar, Spectroscopic characterization of 1,2,4,5-tetracyanobenzene, *Chemical Physics Letters* 460 (2008) 18–22.
- [12] M. J. Frisch, G. W. Trucks, H. B. Schlegel et al., “Gaussian 09, R. A gaussian. Inc., Wallingford CT,” vol. 121, pp. 150–166, 2009. 10.1016/j.cplett.2008.05.037.
- [13] P. B. Nagabalasubramanian, S. Periandym; FT IR and FT Raman, molecular geometry, vibrational assignments, ab initio and density functional theory calculations for 1, 5-methylnaphthalene; *Spectrochimica Acta Part A* 77 (2010) 1099–1107. <https://doi.org/10.1016/j.saa.2010.09.002>
- [14] E. D. Glendening, A. E. Reed, J. E. Carpenter, F. Weinhold, and N. B. O. Version, 3.1 Program Manual, University of Wisconsin, Madison, 1998.
- [15] J. Foresman and E. Frish, Exploring Chemistry with Electronic Structure Methods, Gaussian Inc., Pittsburg, USA, 1996.
- [16] I Fleming, *Frontier Orbitals and Organic Chemical Reactions*, John Wiley and Sons, NewYork, 05–27, (1976).
- [17] Wilson E B, Decius J C & Cross P C, *Molecular vibration: the theory of infrared and Raman vibrational spectra*, Dover Publ Inc New York, 1980. urn: lcp: molecularvibrati00wils: lcpdf: e3fe33b0-1929-46a2-920c-91471ef13761.
- [18] T. Prabhu, S. Periandy, S. Mohan, Spectroscopic (FT IR and FT Raman) analysis and vibrational study on 2, 3-dimethylnaphthalene using ab-initio HF and DFT calculations, *Spectrochimica Acta Part A* 78 (2011) 566–574. <https://doi.org/10.1016/j.saa.2010.11.024>
- [19] V. Krishnakumar, N. Prabavathi, S. Muthunatesan, Density functional theory study of vibrational spectra, and assignment of fundamental vibrational modes of 1-bromo-4-fluoronaphthalene, *Spectrochimica Acta Part A* 70 (2008) 991–996. <https://doi.org/10.1016/j.saa.2007.07.063>
- [20] G. Keresztury, Raman spectroscopy: theory, in: J. M. Chalmers, P. R. Griffiths (Eds.), *Handbook of Vibrational Spectroscopy*, vol. 1, John Wiley & Sons Ltd., 2002, p.71. PrintISBN: 9783527297825| Online ISBN: 9783527602308 |DOI: 10.1002/3527602305.
- [21] L. Goodman, A. G. Ozkabak, S. N. Thakur, Abenchmark vibrational potential surface: ground-state benzene, *J. Phys. Chem.* 95 (1991) 9044–9058. <https://doi.org/10.1021/J100176A008>
- [22] The official web site of National Institute of Advanced Industrial Science and Technology (AIST), Research Information Database (RIO-DB). <http://riodb.ibase.aist.go.jp/riohomee.html>.
- [23] S. George, *Infrared and Raman Characteristic Group Frequencies—Tables and Charts*, thirded., (Wiley, New York, 2001).
- [24] V. Krishnakumar, N. Surumbarkuzhali, S. Muthunatesan, Scaled quantum chemical studies on the vibrational spectra of 4-bromobenzonitrile, *Spectrochimica Acta Part A* 71 (2009) 1810–1813. <https://doi.org/10.1016/j.saa.2008.06.037>
- [25] M Alcolea Palafox and V K Rastogi, Spectra and structure of benzonitriles and some of its simple derivatives, *Asian Journal of Physics* Vol. 22, No 3 (2013) 1-30.
- [26] N. Sundaraganesan, G Elango, S Sebastian and P Subramani, Molecular structure; vibrational spectroscopic studies and analysis of 2-fluoro-5-methylbenzonitrile, *Indian J PURE & APPL PHYS*, Vol 47, July 2009.